

Map Source: lowa Department of Natural Resources, "Natural Resources Geographic Information Systems Library," http://www.igsb.uiowa.edu/nrgislibx/.



Bioregional Context

Julia Badenhope, Matthew Gordy, Colby Fangman, Riley Dunn

Iowa State University | Trees Forever | Iowa Department of Transportation

Historical Vegetation

The vegetation information shown here is derived from township maps made by the General Land Office (GLO) surveys beginning in 1836 through 1859. The vegetation information was digitized in 1996 as a resource for natural resource management and is useful "...for the study of long term ecological processes and as baseline data for the study of present day communities."1

The plant communities mapped by the GLO surveyors varied in classification and the terminology from the original maps has been preserved.

The vegetation types are defined²:

- 1. <u>Field</u>: Cultivated lands of early pioneers.
- 2. <u>Grove</u>: Isolated dense young stand of trees.
- 3. <u>Marsh</u>: Perennial wetlands, basins of irregular shape.
- 4. <u>Prairie</u>: Dominated by prairie grasses with individual or few scattered trees.
- 5.<u>Thicket</u>: Impenetrable blocks of young trees, often thorny.
- 6.<u>Timber</u>: Contiguous blocks of trees extending to the horizon in at least one direction.
- 7. <u>Slough</u>: Like marsh but more linear in shape.

Michael Charles Miller, "Analysis of historic vegetation patterns in Iowa using Government Land Office surveys and a Geographic Information System" (master's thesis, Iowa State University, 1995), 134-135.



J.E. Ebinger, "Presettlement Vegetation of Coles County, Illinois," Transactions of the Illinois Academy of Science (1987): 15-24, quoted in Michael Charles Miller, "Analysis of historic vegetation patterns in Iowa using Government Land Office surveys and a Geographic Information System" (master's thesis, Iowa State University, 1995), 8.